

Title : will be set by the publisher
 Editors : will be set by the publisher
 EAS Publications Series, Vol. ?, 2013

DENSITOMETRY AND THERMOMETRY OF STARBURST GALAXIES

Jeffrey G. Mangum¹, Jeremy Darling², Karl M. Menten³, Christian Henkel³ and Meredith MacGregor^{1, 4}

Abstract. With a goal toward deriving the physical conditions in external galaxies, we present a survey of formaldehyde (H_2CO) and ammonia (NH_3) emission and absorption in a sample of starburst galaxies using the Green Bank Telescope. By extending well-established techniques used to derive the spatial density in star formation regions in our own Galaxy, we show how the relative intensity of the $1_{10} - 1_{11}$ and $2_{11} - 2_{12}$ K-doublet transitions of H_2CO can provide an accurate densitometer for the active star formation environments found in starburst galaxies (*cf.* Mangum *et al.* 2008). Similarly, we employ the well-established technique of using the relative intensities of the (1,1), (2,2), and (4,4) transitions of NH_3 to derive the kinetic temperature in starburst galaxies. Our measurements of the kinetic temperature constrained spatial density in our starburst galaxy sample represent the first mean density *measurements* made toward starburst galaxies. We note a disparity between kinetic temperature measurements derived assuming direct coupling to dust and those derived from our NH_3 measurements which points to the absolute need for direct gas kinetic temperature measurements using an appropriate molecular probe. Finally, our spatial density measurements point to a rough constancy to the spatial density ($10^{4.5}$ to $10^{5.5} \text{ cm}^{-3}$) in our starburst galaxy sample. This implies that the Schmidt-Kennicutt relation between L_{IR} and M_{dense} : (1) Is a measure of the dense gas mass reservoir available to form stars, and (2) Is not directly dependent upon a higher average density driving the star formation process in the most luminous starburst galaxies.

¹ National Radio Astronomy Observatory, 520 Edgemont Road, Charlottesville, VA 22903-2475, USA; e-mail: jmangum@nrao.edu

² Center for Astrophysics and Space Astronomy, Department of Astrophysical and Planetary Sciences, Box 389, University of Colorado, Boulder, CO 80309-0389, USA; e-mail: jdarling@origins.colorado.edu

³ Max Planck Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany; e-mail: kmenten@mpifr-bonn.mpg.de, chenkel@mpifr-bonn.mpg.de

⁴ Harvard University, Cambridge, MA 02138, USA; e-mail: mmacgreg@fas.harvard.edu

1 Introduction

Studies of the distribution of carbon monoxide (CO) emission in external galaxies (*cf.* Young & Scoville 1991) have pointed to the presence of large quantities of molecular material in these objects. These studies have yielded a detailed picture of the molecular mass in many external galaxies. But, because emission from the abundant CO molecule is generally dominated by radiative transfer effects, such as high optical depth, it is not a reliable monitor of the physical conditions, such as spatial density and kinetic temperature, quantities necessary to assess the potential for star formation. Emission from less-abundant, higher-dipole moment molecules are better-suited to the task of deriving the spatial density and kinetic temperature of the dense gas in our and external galaxies. For this reason, emission line studies from a variety of molecules have been made toward mainly nearby galaxies (see Henkel *et al.* 1991 for a review).

Formaldehyde (H_2CO) has proven to be a reliable density and kinetic temperature probe in Galactic molecular clouds. Existing measurements of the H_2CO $1_{10} - 1_{11}$ and $2_{11} - 2_{12}$ emission in a variety of galaxies by Baan *et al.* (1986, 1990, 1993) and Araya *et al.* (2004) have mainly concentrated on measurements of the $1_{10} - 1_{11}$ transition. One of the goals of the present study was to obtain a uniform set of measurements of both K-doublet transitions with which the physical conditions, specifically the spatial density, in the extragalactic context could be derived. Using the unique density selectivity of the K-doublet transitions of H_2CO within the framework of a radiative transfer model which incorporates the Large Velocity Gradient (LVG) approximation, we have measured the spatial density in a sample of galaxies exhibiting starburst phenomena and/or high infrared luminosity. Our ammonia (NH_3) measurements of these same galaxies, together with our LVG model, provide kinetic temperature measurements which can further constrain the range of allowed densities derived from our H_2CO measurements.

2 Galaxy Sample Summary

The galaxy selection criteria and summary measurement results are as follows: Selection Criteria: $\text{Dec}(\text{J2000}) > -40^\circ$; $L(60\ \mu\text{m}) > 50\ \text{Jy}$ and/or bright HCN and CO emission. H_2CO measurements summary: GBT $\theta_b = 51\ \text{arcsec}$ ($\lambda = 2\ \text{cm}$) and $153\ \text{arcsec}$ ($\lambda = 6\ \text{cm}$); 26 of 56 galaxies detected in at least one H_2CO transition; $1_{10} - 1_{11}$ emission/absorption = 6/19; $2_{11} - 2_{12}$ emission/absorption = 1/13; 17 of 26 detections are new extragalactic discoveries of H_2CO . Figure 1 shows two examples of our H_2CO results. Emission in either of the two H_2CO transitions in this study is a sign of high spatial density. Observed emission in the H_2CO $2_{11} - 2_{12}$ transition toward Arp 220 signals a relatively high density ($\log(n(\text{H}_2)(\text{cm}^{-3})) \gtrsim 5.0$). Absorption in both H_2CO transitions toward NGC 253, on the other hand, signals an average density that is lower ($\log(n(\text{H}_2)(\text{cm}^{-3})) \lesssim 5.0$).

NH_3 measurements summary: GBT $\theta_b = 30\ (\lambda = 1.3\ \text{cm})\ \text{arcsec}$; 12 of 19 galaxies detected in at least one NH_3 transition; NH_3 emission/absorption = 7/5; 4 of 12 detections are new extragalactic discoveries of NH_3 . Figure 1 shows two

examples of our NH_3 measurement results. As these examples reveal, our NH_3 measurements reveal both emission and absorption in the (1,1), (2,2), (4,4), and absorption in the (2,1) transition. *Emission* in the NH_3 (2,1) transition is a hallmark of high infrared intensity (*cf.* Morris *et al.* 1973). NH_3 (2,1) absorption, on the other hand, requires only a strong background continuum source.

3 Density and Kinetic Temperature Measurements

For the H_2CO sources in which we have detected both the $1_{10} - 1_{11}$ and $2_{11} - 2_{12}$ transitions we have applied our technique to derive the spatial density assuming kinetic temperatures derived both from dust emission and our NH_3 measurements (Table 1). Noteworthy are the generally higher, and in a few cases *much* higher, kinetic temperatures derived using our NH_3 measurements. In many cases these high kinetic temperatures have been previously noted (*cf.* Ott *et al.* 2005), pointing to the importance of a direct measurement of the kinetic temperature in these starburst environments.

4 Conclusions

The H_2CO and NH_3 measurements presented allow us to derive the kinetic temperature and spatial density in our starburst galaxy sample, representing the first mean density *measurements* made toward starburst galaxies. The disparity between kinetic temperature measurements derived assuming direct coupling to dust and those derived from our NH_3 measurements points to the absolute need for direct gas kinetic temperature measurements using an appropriate molecular probe. Finally, our spatial density measurements point to a rough constancy to the spatial density ($10^{4.5}$ to $10^{5.5} \text{ cm}^{-3}$) in our starburst galaxy sample. This implies that the Schmidt-Kennicutt relation between L_{IR} and M_{dense} : (1) Is a measure of the dense gas mass reservoir available to form stars, and (2) Is not directly dependent upon a higher average density driving the star formation process in the most luminous starburst galaxies.

References

- Araya, E., Baan, W. A., & Hofner, P. 2004, ApJS, 154, 541
- Baan, W. A., Güsten, R., & Haschick, A. D. 1986, ApJ, 305, 830
- Baan, W. A., Henkel, C., Schilke, P., Mauersberger, R., & Güsten, R. 1990, ApJ, 353, 132
- Baan, W. A., Haschick, A. D., & Uglesich, R. 1993, ApJ, 415, 140
- Henkel, C. Baan, W. A., & Mauersberger, R 1991, A&AR, 3, 47
- Mangum, J. G., Darling, J., Menten, K. M., & Henkel, C. 2008, ApJ, 673, 832
- Morris, M., Zuckerman, B., Palmer, P., & Turner, B. E. 1973, ApJ, 186, 501
- Ott, J., Weiss, A., Henkel, C., & Walter, F. 2005, ApJ, 629, 767
- Young, J. S. & Scoville, N. Z. 1991, ARA&A, 29, 581

Table 1. Derived Kinetic Temperatures and Spatial Densities

Galaxy	α (J2000)	δ (J2000)	v_{hel} (km s ⁻¹)	T_D, T_K (K)	$\log(n(H_2))$ (cm ⁻³)
NGC 253	00:47:33.1	-25:17:18	251	34 (D) 81 $\pm$ 2	5.08 $\pm$ 0.02 5.00 $\pm$ 0.03
IC 342	03:46:49.7	+68:05:45	31	30 (D) 38 $\pm$ 3 104 $\pm$ 7	5.05 $\pm$ 0.11 5.05 $\pm$ 0.11 4.89 $\pm$ 0.14
M 82	09:55:52.2	+69:40:47	203	45 (D) 58 $\pm$ 16	4.95 $\pm$ 0.02 4.92 $\pm$ 0.05
NGC 3079C1	10:01:57.8	+55:40:47	1150	32 (D) > 260	5.56 $\pm$ 0.02 < 4.42
NGC 3079C2				32 > 180	5.47 $\pm$ 0.02 < 4.94
IC 860	13:15:04.1	+24:37:01	3866	40 (D) > 52	5.70 $\pm$ 0.02 < 5.67
M 83	13:37:00.9	-29:51:57	518	31 (D) 48 $\pm$ 9	5.38 $\pm$ 0.08 5.39 $\pm$ 0.09
Arp 220	15:34:57.1	+23:30:11	5434	44 (D) > 180	5.64 $\pm$ 0.02 < 5.00
NGC 6946	20:34:52.3	+60:09:14	48	30 (D) 44 $\pm$ 7	5.05 $\pm$ 0.20 5.06 $\pm$ 0.21

D indicates dust temperature (see Mangum *et al.* 2008).




Fig. 1. Top Row: H₂CO 1₁₀–1₁₁ (top) and 2₁₁–2₁₂ (bottom) spectra from Arp 220 (left) and NGC 253 (right). Bottom Row: NH₃ (4,4), (2,2), (1,1), and (2,1) (top-to-bottom) spectra from IC 342 (left) and NGC 660 (right).